

Sumontha. Institut Royal des Sciences Naturelles de Belgique, Brussels (IRSNB 17014). Verified by Georges Lenglet (IRSNB). First provincial record. The species is thus presently recorded in Thailand from Kanchanaburi, Nakhon Ratchasima, Narathiwat, Phang-Nga, Phuket, Ranong, Trang, and Trat provinces (Chandler et al. 1999. Amphibians and Reptiles of Peninsular Malaysia and Thailand. An Illustrated Checklist. Bushmaster Publications, Würselen, 240 pp.; Nabhitabhata et al., "2000" 2004. Checklist of Amphibians and Reptiles in Thailand, Office of Environmental Policy and Planning, Bangkok; Pauwels and Bauer 2001. Herpetol. Rev. 32:119).

Submitted by **OLIVIER S. G. PAUWELS**, Department of Recent Vertebrates, Institut Royal des Sciences naturelles de Belgique, Rue Vautier 29, 1000 Brussels, Belgium (e-mail: osgpauwels@yahoo.fr); and **MONTRI SUMONTHA**, Ranong Marine Fisheries Station, 157 M. 1, Saphan-Pla Rd., Paknam, Muang, Ranong 85000, Thailand (e-mail: knotsnake2211@yahoo.com).

**HOLASPIS GUENTHERI** (Günther's Gliding Lizard). EQUATORIAL GUINEA: Monte Alen National Park. July 2001. T. Stévant. Institut Royal des Sciences Naturelles de Belgique, Brussels (IRSNB 16638). Verified by Georges L. Lenglet (IRSNB). Adult specimen found active by day in secondary forest nr. ECOFAC station and hotel (01°39'N, 10°18'E, ca. 800 m elev.). New record for Park and for Centro Sur Province (Lasso et al. 2002. Graellsia 58[2]:21–34; De la Riva 2004. Quercus 215:56–61). We thank Ignacio J. De la Riva (Museo Nacional de Ciencias Naturales, Madrid) for comments. This is Contribution 89 of the Smithsonian Institution's Gabon Biodiversity Program.

Submitted by **OLIVIER S. G. PAUWELS**, Smithsonian Institution, National Zoological Park, Monitoring and Assessment of Biodiversity Program, Gamba, Gabon; mailing address: Département des Vertébrés Récents, Institut Royal des Sciences naturelles de Belgique, Rue Vautier 29, 1000 Brussels, Belgium (e-mail: osgpauwels@yahoo.fr); and **TARIQ STÉVART**, Missouri Botanical Garden, mailing address: Laboratoire de Botanique systématique et de Phytosociologie, Université Libre de Bruxelles, CP 169, Av. F. D. Roosevelt 50, 1050 Brussels, Belgium (e-mail: tstevart@yahoo.com).

**LIOLAEMUS TELSEN** (NCN). ARGENTINA: RIO NEGRO: 9 DE JULIO DEPARTMENT: Ruta Provincial 60, 65.6 km S Police Post El Rincón, near Cerro Corona (41°23'40.8"S 66°57'33.3"W) 1425 m elev. 2 November 2006. L. J. Avila. Museo de La Plata herpetological collection (MLP.S 2596). Verified by J. A. Sclaro. Previously known only from type locality—"Chubut, Argentina (some 80 km to west from Telsen)." First province record and northernmost record for the species, extending the known distribution ca. 100 km NE from the type locality in Chubut Province (Cei and Sclaro 1999. Revue Française de Aquariologie 1–2:79–82).

Submitted by **NICOLAS FRUTOS**, **CRISTIAN H. F. PEREZ**, **MONICA KOZYKARISKI**, **MARIANA MORANDO**, and **LUCIANO JAVIER AVILA**, CENPAT-CONICET, Boulevard Almirante Brown s/n, U9120ACV, Puerto Madryn (Chubut), Argentina (e-mail: avila@cenpat.edu.ar).

**PHELSUMA MADAGASCARIENSIS GRANDIS** (Madagascar

Giant Day Gecko). USA: FLORIDA: MONROE Co.: Vaca Key, Marathon, 1117 28th Street (24°42'26.924"N, 81°5'52.051"W, NAD83; elev. <1 m). 15 October 2006. Richard S. Lake. Verified by Josiah H. Townsend. Florida Museum of Natural History (photographic vouchers UF 150102–04). New island record and extends the range 16.5 km SW of the closest known locality on Grassy Key (Krysko et al. 2003. Florida Sci. 66:222–225; Krysko and Sheehy 2005. Carib. J. Sci. 41:169–172). Neonate and adults observed on side of house. On 27 October 2006, three neonates (photographic voucher UF 150105) and three adults were observed at this site. On 7 January 2007, several individuals were again observed at this site, as well as an adult (photographic voucher UF 150187) at 1313 28th Street (24°42'25.056"N, 81°5'51.54"W, NAD83; elev. <1 m).

Submitted by **KENNETH L. KRYSKO**, Division of Herpetology, Florida Museum of Natural History, Dickinson Hall, University of Florida, Gainesville, Florida 32611, USA (e-mail: kennedyk@flmnh.ufl.edu); **RICHARD S. LAKE**, 302 Daniel Paul Drive, Archdale, North Carolina 27263, USA; and **CARL D. MAY**, 1311 N Lakeside Drive, Lake Worth, Florida 33460, USA (e-mail: cdmfla@aol.com).

**PHRYNOSOMA DOUGLASII** (Pygmy Short-horned Lizard). USA: WYOMING: LINCOLN Co.: T21N, R113W, SW 1/4 Section 16. 08 August 1994. Kris H. Johnson. YPM 10176. Juvenile, with distinct umbilical scar. Verified by Greg Watkins-Colwell. New species record for Wyoming, and new county record for the genus (Baxter and Stone 1985. Amphibians and Reptiles of Wyoming, Wyoming Game and Fish Dept. Bull. 16; Cerovski et al. 2004. Atlas of Birds, Mammals, Amphibians, and Reptiles in Wyoming, Wyoming Game and Fish Dept.). Extends known range ca. 70 km E of the Idaho/Wyoming border (e.g., Hodges 2003. Digital Morphology, accessed 18 December 2006 at [http://digimorph.org/specimens/Phrynosoma\\_douglasii/](http://digimorph.org/specimens/Phrynosoma_douglasii/); Stebbins 2003. A Field Guide to Western Reptiles and Amphibians, 3<sup>rd</sup> Ed., Houghton Mifflin Co.).

Submitted by **KRIS H. JOHNSON**, U.S. Fish and Wildlife Service Ecological Services Division, 620 East Greene Street, Carlsbad, New Mexico 88220, USA.

**SCELOPORUS CYANOGENYS** (Blue Spiny Lizard). USA: TEXAS: NUECES Co.: South of Bishop, ca. 20 m S of County Road 4, 1 road km E of business Highway 77 (27°34.130N, 097°48.358'W). 08 October 2006. Randy L. Powell. Verified by Travis J. LaDuc. Texas Natural History Collections (TNHC 65743). New county record, extends range east one county (Dixon 2000. Amphibians and Reptiles of Texas, 2nd ed. Texas A&M Univ. Press, College Station. 421 pp.).

Submitted by **RANDY L. POWELL** Department of Biological and Health Sciences, MSC 158, Texas A&M University, Kingsville, Texas 78363, USA; e-mail: randy.powell@tamuk.edu.

**SCINCELLA GEMMINGERI** (Forest Ground Skink) MÉXICO: QUERÉTARO: MUNICIPALITY OF CADEREYTA DE MONTES: 0.5 km E de Maconi. (20°49'48"N, 99°32'99"W), 1920 m elev. 23 May 1996. Laura Contreras Narváez. Verified by Luis Canseco Márquez. Colección Herpetológica, Facultad de Ciencias, UNAM (MZFC 9781). First record for the municipality and second record for the state, extending the known distribution 44.6 km SW from 1.6 km

W Landa de Matamoros in the Sierra Gorda region, Querétaro (Ketchersid 1974. Herpetofauna of Two Biogeographic Transects in Eastern Mexico. Unpublished dissertation. Texas A&M Univ., College Station), and ca. 40.9 km range extension W from the closest known locality at La Placita, Hidalgo (García-Vázquez 2004. Revisión Taxonómica del Genero *Scincella* [Lacertilia: Scincidae] de México. Unpublished thesis, Univ. Autón. de Puebla, Puebla). The specimen was caught in riparian vegetation.

Submitted by **URI OMAR GARCÍA-VÁZQUEZ** and **ANDRÉS ALBERTO MENDOZA-HERNÁNDEZ**, Museo de Zoología, Facultad de Ciencias, UNAM, A.P. 70-399, México D. F. 04510 (e-mail: urigarcia@gmail.com).

## SERPENTES

**AHAETULLA FASCIOLATA** (Speckled-head Whip Snake). THAILAND: RANONG PROVINCE: Muang District: forest stream near Klong Hat Som Paen (ca. 9°57'N, 98°41'E). 21 February 2004. M. Sumontha. Queen Saovabha Memorial Institute, Bangkok (QSMI 600). Verified by L. Chanhome (QSMI). Found on a tree along a forest stream in syntopy with *Ahaetulla prasina* (voucher at the Institut Royal des Sciences Naturelles de Belgique, Brussels: IRSNB 16997). First provincial record and extension of ca. 175 km N from the previous northernmost locality which is situated at Raman Forest Park, Takua Thung District, Phang-Nga Province (Pauwels et al. 2002. Nat. Hist. J. Chulalongkorn Univ. 2[1]:25–30). We thank Lawan Chanhome (QSMI) and Wudtichai Wungkahart (Ranong Marine Fisheries Station) for working facilities.

Submitted by **OLIVIER S. G. PAUWELS**, Department of Recent Vertebrates, Institut Royal des Sciences Naturelles de Belgique, Rue Vautier 29, 1000 Brussels, Belgium (e-mail: osgpauwels@yahoo.fr); and **MONTRI SUMONTHA**, Ranong Marine Fisheries Station, 157 M. 1, Saphan-Pla Rd., Paknam, Muang, Ranong 85000, Thailand (e-mail: knotsnake2211@yahoo.com).

**CLONOPHIS KIRTLANDII** (Kirtland's Snake). USA: MISSOURI: CLARK CO.: Near Wayland (T65N, R6W). 28 April 2006. Christopher Shulse. Verified by J. M. Jones and R. Daniel. Dean E. Metter Memorial Collection, University of Missouri, Columbia (UMC 7936). Rediscovery in state, new county record, only second report west of Mississippi River, and extends the range in Missouri by ca. 51 km. One dead specimen was found on the shore of a constructed pond. Four live specimens were observed near the pond from 28 April – 31 May 2006. This species was first recorded in Missouri in 1964. A single live female was collected on 10 May 1964 in Marion County near Taylor, Missouri (Jones 1967. Herpetologica 23:66–67). No additional specimens were observed until those described herein. Therefore, this species was not included as part of Missouri's herpetofauna by Johnson (2000, The Amphibians and Reptiles of Missouri. Second Ed. Missouri Dept. Conserv., Jefferson City. 400 pp.). Based on these discoveries, *C. kirtlandii* is now considered extant in Missouri and is state ranked as critically imperiled (Missouri Department of Conservation 2007. Missouri Species and Communities of Conservation Concern. Missouri Dept. Conserv., Jefferson City. 50 pp.) This species is known to occur in six other states where it is listed as

either state endangered or threatened (Ernst and Ernst 2003. Snakes of the United States and Canada. Smithsonian Inst. Press, Washington, DC. 668 pp.), and has an imperiled global rank (Missouri Department of Conservation 2007, *op. cit.*). Additional studies are planned to investigate its distribution and abundance in northeast Missouri (J. Briggler, pers. comm.).

Submitted by **CHRISTOPHER SHULSE**, 212 Tucker Hall, Division of Biological Sciences, University of Missouri, Columbia, Missouri 65211, USA; e-mail: cdsxzc@mizzou.edu.

**CROTALUS WILLARDI WILLARDI** (Arizona Ridge-nosed Rattlesnake). MÉXICO: SONORA: MUNICIPIO DE NOGALES: Sierra de Pinitos, unnamed canyon on private land S of Los Picos Peaks, ca. 20 km SE (straight line) from the city of Nogales (ca. 31°13'10.15"N, 110°49'30.04"W; UTM: 518000E, 3450000N; horizontal datum ITRF92; ellipsoid GRS80; UTM zone 12), ca. 1524 m elev. 05 May 2006. Robert Villa, Paul Condon, and Trevor Hare. Verified by George L. Bradley. UAZ Sonoran Herpetological Photographic Voucher Initiative, UAZ 56499-PSV (skin with rattle). Westernmost locality for the species in México, extending known range ca. 50 km NW (straight line) from Cananea, Sonora (UAZ 27945). This is also the only record west of the Santa Cruz River. The snake was found in Madrean evergreen woodland.

Submitted by **ROBERT A. VILLA**, 1125 East Missouri Street, Tucson, Arizona 85714, USA (e-mail: herpsandviolin@aol.com); **PAUL T. CONDON**, 8832 East Buckboard Road, Tucson, Arizona 85749, USA; **TREVOR A. HARE** and **SERGIO AVILA-VILLEGAS**, Sky Island Alliance, 738 North 5th Avenue, Suite 201, Tucson, Arizona 85705, USA; and **DAVID G. BARKER**, Vida Preciosa International, Inc., P.O. Box 300, Boerne, Texas 78006, USA.

**DENDRELAPHIS KOPSTEINI** (Kopstein's Bronzeback Tree Snake). BRUNEI DARUSSALAM: BELAIT DISTRICT: Sungei Rampayoh (04°26'N, 114°24'E). Earlier reported as *Dendrelaphis formosus* (Das 1999. Malayan Nat. J. 53[4]:349–350; Das 2007. A Pocket Guide. Amphibians and Reptiles of Brunei. Natural History Publications [Borneo] Sdn Bhd. Kota Kinabalu. viii + 200 pp.), this species was recently described as new on the basis of specimens formerly identified as *D. formosus* from Thailand, Peninsular Malaysia, Singapore, and Sumatra (Vogel and van Rooijen 2007. Zootaxa 1394:25–45) and diagnosable from congeners in showing the following combination of characters: dark stripe covers only the lower part of temporal region, terminating at rear of neck; nuchal region red in life when inflated; eyes large; two supralabials in contact with orbit of eye. First record for Borneo.

Submitted by **INDRANEIL DAS**, Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia; e-mail: idas@ibec.unimas.my.

**LEPTODEIRA SPLENDIDA EPHIPPIATA** (Saddled Cat-eyed Snake). MÉXICO: SONORA: Municipio de Yécora, 45.7 km (by Mex. Hwy. 16) E Yécora (28.40430°N, 108.67572°W), 1280 m elev. 06 July 2006. E. Enderson and R. Bezy. UAZ 56548-PSV. Verified by George Bradley. Northernmost record for Sonora, extending the range ca. 232 km (airline) N from the vicinity of Agua Marín, Sonora (Duellman 1958. Bull. Amer. Mus. Nat. Hist. 114:1–